
STICKING PROBABILITY OF H2 AND HD ON NOBLE METAL SURFACES(U) CORNELL UNIV ITHACA NY LAB OF ATOMIC AND SOLID STATE PHYSICS M D STILES ET AL. 1980 TR-6 N00014-80-C-0489 F/G 7/4 UNCLASSIFIED NL

1/1

AD-8147 061





OFFICE OF NAVAL RESEARCH

Research Contract No. NOO014-80-C-0489

TECHNICAL REPORT No. 6

STICKING PROBABILITY OF H2 AND HD ON NOBLE METAL SURFACES

by

M.D. Stiles and J.W. Wilkins

Prepared for Publication

in

Physical Review Letters

Laboratory of Atomic and Solid State Physics Cornell University Ithaca, NY 14853


Reproduction in whole or part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited.

OTIC FILE CUPY



84 10 26 017



Sticking Probability of ${\rm H_2}$ and ${\rm HD}$ on Noble Metal Surfaces

M. D. Stiles and J. W. Wilkins

Laboratory of Atomic and Solid State Physics Cornell University Ithaca, NY 14853

One-phonon distorted-wave Born approximation calculations of the energy-dependent phonon-assisted sticking coefficients of $\rm H_2$ and $\rm HD$ at smooth metal surfaces yield peaks in the sticking coefficient as a function of energy due to selective adsorption resonances, the peaks being strong for $\rm HD$ and weak for $\rm H_2$. Furthermore the calculation does not explain the observed dependence of the sticking coefficient of $\rm H_2$ on the rotational populations.

PACS numbers: 82.65.My., 68.30.+z.

FLLMED

7

11-84

DTIC